
Trauma-Informed Care is a
Culture that Helps Us All

Lea C. Watson, MD, MPH Visiting Professor of Psychiatry and Geriatrics University of Colorado School of Medicine



1




2



"Trauma pervades our culture, from personal functioning through social relationships, parenting, education, popular culture, economics, and politics. In fact, someone *without* the marks of trauma would be an outlier in our society."

~Dr. Gabor Mate in The Myth of Normal



SHOUT OUT!

This talk includes content from a 3-hour workshop presented at the AMDA Annual Conference March 2023 Tampa, FL Creating a Sustainable Trauma-Informed Care Culture for Residents and Staff Post Pandemic

"The health care system is populated by trauma survivors, both those providing and receiving care."

(Fleishman, 2019)

Paige Hector, LMSW, Lea Watson, MD Lisa Lind, PhD, Allison Villegas, PA-C


5

Emotional and Psychological Trauma

"Result of **extraordinarily stressful events** that shatter your sense of security, making you feel **helpless** in a dangerous world. Often involve a threat to life or safety, but any situation that leaves you **feeling overwhelmed** and **isolated** can result in trauma, even if it doesn't involve physical harm. The more **frightened and helpless** you feel, the more likely you are to be traumatized."

(emphasis added)

Emotional and Psychological Trauma https://www.helpguide.org/articles/ptsd-trauma/coping-with-emotional-and-psychological-trauma.htm



Emotional Exhaustion Among Health Care Workers (HCWs)

- 40% of nurses and 23.8% of physicians plan to exit their practice in the next 2 years
- Comparison of post 9/11 combat veterans to HCWs during the pandemic shows equivalent rates of moral injury in both groups
 - Emotional exhaustion rates among HCWs were already considered alarmingly high before the pandemic

"Emotional exhaustion is a chronic state of physical and emotional depletion that results from excessive job demands and continuous hassles." (Psychology Wiki)

JAMA Network Open. 2022;5(9):e2232748. doi:10.1001/jamanetworkopen.2022.32748

8



Losses Related to Aging and Illness

- Independence living space, driving
- Daily living skills (ADLs and IADLs)
- Finances
- Death of partner or spouse
- Loss of meaningful roles
- Health and cognition
- Nursing home "placement"

Sources of Medical Trauma

- Interactions with 'the system'
- $\bullet \;\;$ Communication that is too technical, too vague, too infrequent or too frequent
- · Medication side effects
- Illness-related symptoms (e.g., pain, shortness of breath, racing heartbeat, Gl distress, physical weakness, difficulty swallowing/choking)
- Loud noises, falls, nightmares
- · IV placement, limited movement, restraints
- Exposure to sounds, lights, odors
- Private areas being seen/touched by multiple people
- Exposure to needles, blood, temperature changes
- Feeling isolated, powerless, vulnerable, depressed
- Fearing for one's wellbeing and life
- Being in the dark
- Being treated or talked to "like a child"

10

Hospitalization can cause trauma

Especially in those living with dementia

Waiting can trigger feelings associated with neglect, abandonment


Fragmented care

Propensity for over-testing

Transfer and transitions = uncertainty, discomfort, overwhelm, fear, anxiety

Goals of care interrupted

11



TIC is


- TIC is person-centered care
- TIC is a fundamental perspective
- TIC is an integrative framework
- TIC is a relational posture towards everyone who is involved
- TIC is a workplace culture

TIC is 'NOT'

- TIC is NOT a training on PTSD
- TIC is NOT based solely on the medical model
- TIC is NOT just a prescribed protocol or set of skills
- TIC is NOT just for residents
- TIC is NOT just for people who have PTSD

Ashley Swinson, MSW, LCS


13





14

"Trauma is a psychic injury, lodged in our nervous system, mind, and body, lasting long past the originating incident(s), triggerable at any moment."

~Dr. Gabor Mate







Healthcare seeking						
Can be a proxy for getting emotional needs met	Creates significant risk for latrogenic harm	Is often confounded by complex medical history	Places high burden on the clinician			

What are Triggers?

- Triggers are reminders of dangerous or frightening things (or people) that happened in the past* and the person experiences the event all over again (even if the current environment is "safe")
- Triggers come without warning and can be ANYTHING
 - Triggers can be puzzling or disturbing for others, especially when the person associates us or something we are doing with trauma
- The person may not even associate the trigger with the event or know it's happening
 - Watch for stiffening, combativeness, crying out, withdrawal, sudden silence, etc.

*The past can be moments ago or many years ago.

20

Triggers (trauma reminders) can be interpreted as... "I'm not safe." "I can't protect myself." "I'm going to die."

lanssen S. Assessing for PTSD in Terminally III Patients. The New Social Worker. Accessed April 29, 2



Behavioral Expressions

- Yelling
- Arguing
- OCD and other anxiety disorders
- Isolation, withdrawal
- Protective gestures
- Aggression (verbal and physical)
- Resistance to care
- Declining care
- \bullet Self injurious coping mechanisms drugs, alcohol, prostitution
- Unwelcome sexual expression

23



Two Key Questions

1. How could this behavior make sense as a reaction to past trauma?

These may be COPING MECHANISMS that

made perfect sense at the time of a

traumatic experience although they may


no longer suit the current circumstance.

"Nor are they character faults; though

they may cause us difficulty now, they began as modes of survival." (Dr. Mate)

2. What might this person need to avoid reliving their trauma in the future?

(Crisis and Trauma Resource Institute



Safety

- **Physical safety** includes the physical plant, security measures, disaster planning, policies and procedures.
- Social safety refers to the ability to be a part of a group, to listen and to be heard, to be able to play a role in conflict resolution, to use one's intelligence and creativity to serve a group process without engaging in behavior or activities that destroy the integrity of the self or the group.
- Moral safety reflects an environment that actively defines and redefines a moral universe of integrity, responsibility, honesty, tolerance, compassion, peace, nonviolence, justice, and an abiding concern for human rights.

The Sanctuary Model: Through the Lens of Moral Safety by Sandra Bloom, https://sandrabloom.com/wgcontent/uploads/2017-BLOOM-THE-SANCTUARY-MODEL-THROUGH-THE-LENS-OF-MORAL-SAFETY.pd

26

Trust and Transparency

Trust – being vulnerable and sharing personal information can feel risky

- Gentle, low-key approach, no 'agenda'
- Confidentiality and privacy are key

Transparency - organizational operations and decisions are transparent

- Predictability with processes and daily activities
- Emphasis is not on "getting it right all the time" but rather how situations are handled when circumstances provoke feelings of being

Tributer and Comment of the Comment

Creating a traumainformed organization is a fluid, ongoing process; it has no completion date.



(SAMHSA, 2014)

28

F699 Trauma-Informed Care

(Guidance issued in 2022)

"The facility must ensure that residents who are trauma survivors receive culturally competent, trauma-informed care in accordance with professional standards of practice and accounting for residents' experiences and preferences in order to eliminate or mitigate triggers that may cause re-traumatization of the resident."

29

F699 Trauma-Informed Care

(topics included in the Guidance to Surveyors)

- 6 principles of trauma
- "Assessment" CMS advises a multi-pronged approach to identifying a resident's history of trauma
- Triggers and retraumatization
- Cultural "competency" defined by CMS
 - A "set of behaviors and attitudes held by clinicians that allows them to communicate effectively with individuals of various cultural backgrounds and to plan for and provide care that is appropriate to the culture and to the individual."

F699 Trauma-Informed Care, cont. (topics included in the Guidance to Surveyors)

- Care planning to minimize or eliminate the effect of the trigger on the resident
- Care planning to address cultural preferences
- Language verbal and written communication (e.g., forms)
 Food preparation and choices
- Clothing
- Physical contact or provision of care by a member of the opposite sex
- Cultural etiquette, e.g., eye contact
 Activities that are culturally relevant
- Religious or spiritual preferences throughout stay and at the end of life
- Monitoring delivery of care and services
 - Do the interventions mitigate or reduce the impact of identified triggers

31



32

Trauma-Informed Climate Scale – 10 Assessing perceptions Measures the extent to which employee rights, freedoms, and contributions are valued within the agency Be clear about the intention with requesting staff to complete this questionnaire: How can you create a sense of safety? How will you uphold trust and transparency?

	KEY PRINCIPLES						
	Safety	Trustworthiness and Transparency	Peer Support	Collaboration and Mutuality	Empowerment, Voice, and Choice	Cultural, Historical, and Gender Issues	
SAMHSA 's	10 IMPLEMEN	10 IMPLEMENTATION DOMAINS					
Concept of Trauma and Guidance for a Trauma-Informed	Governance and Leadership	- How does agency leadership communicate its support and guidance for implementing a trauma-informed approach? - How do the agency's mission statement and/or written policies and procedures include a commitment to providing trauma-informed services and supports? - How do leadership and governance structures demonstrate support for the voice and participation of people using their services with how trauma histories?					
Approach GOAL: Stimulate change-focused discussion	Policy	 How do the agency's written policies and procedures include a focus on trauma and issues of safety and confidentiality? How do the agency's written policies and procedures recognize the pervissiveness of trauma in the lives of people using services, and express a commitment for reducing re-traumatization and promoting well-being and recovery. How do the agency's staffing policies demonstrate a commitment to staff training on providing services and supports that are culturally relevant and trauma-informed as part of staff orientation and in-service training? 					
https://store.samhsa.gov/s ma14-4884.pdf	system/files/s	How do human resexperienced traum What policies and services and peer governance policy.	a? procedures are in p supports in meanir	place for including tr ngful and significant	auma survivors/pec	ple receiving	

- DO respond with validating language. For example, "I'm really glad you told me this will help us take the best possible care of you."
 DON'T try to investigate or ask for details right away allow them to talk.
- "• If they are getting upper or going into disturbing material, gently close the conversation and follow up with a clinical referral right away
 DO document any reported traumas and inform the clinical team. Include all known or
 suspected trauma triggers associated with the disclosed experience. This helps the team
- DO let the resident know that you will need to let a few key staff members know about "what happened" so that staff can avoid doing things that trigger difficult memories.
- Do refer to the disclosed experience in general terms. Avoid naming "what happened" unless the resident defines it in a given way.
- DO let the resident know that they won't need to talk about "what happened" if they
 don't want to but they may find that they do want to talk about it as time goes on. Let the resident know someone can be available for them to talk to if and when they are ready, including right away. Follow up.

 DO uphold the resident's privacy, even if the information is unusual.

 DO assess current safety. Was it a recent event or far in the past?

 Barbara L. G.
 Barbara L. G.
 Barbara L. G.

35

Primary Care PTSD Screen for DSM-5 (PC-PTSD-5) In the past month, have you ... 1. had nightmares about the event(s) or thought about the event(s) YES when you did not want to? 2. tried hard not to think about the event(s) or went out of your way to YES NO avoid situations that reminded you of the event(s)? 3. been constantly on guard, watchful, or easily startled? 4. felt numb or detached from people, activities, or your surroundings? YES NO 5. felt guilty or unable to stop blaming yourself of others for the event(s) or any problems the events may have caused? NO Total score is sum of "YES" responses in items 1-5.

Indirect Screening

- We can always be engaged in indirect screening.

 Especially for residents with cognitive impairment and for residents who do not wish to engage in direct screening.
- During intake and day-to-day care, pay attention to comments/actions that could indicate symptoms of traumatic stress.
- After sufficient trust has been established, ask permission to discuss observations.
- If discussion indicates presence of symptoms of traumatic stress, ask if they want to speak to someone. If so, make a referral.
- In the plan of care, identify all potential trauma symptoms and triggers, as well as

37

Universal **Precautions** Model

Gloving and handwashing no matter the hazard level

Assume all individuals have a history of trauma and glove up metaphorically to reduce possibility of triggering or re-traumatizing others.



rauma-Informed Organization Change Manual, <a href="http://socialwork.buffalo.inters/institute-on-trauma-and-trauma-informed-care/Trauma-Informed-care/Trauma-care/Trauma-Informed-care/Trauma-Informed-care/Trauma-Informed-care/Trauma-Informe

38

Trauma-Informed

CARE